GLOUCESTER CO. SPECIAL SERVICES SCHOOL DISTRICT NONPUBLIC NURSING PROGRAM 1340 Tanyard Road Sewell, NJ 08080 (856) 468-6530 x1045

ST. MARGARET REGIONAL SCHOOL 773 Third Street Woodbury Heights, NJ 08097 (856) 845-5200 Fax (856) 845-2405

PRESCRIBED MEDICATION ADMINISTRATION IN SCHOOL DISTRICT REGULATIONS REGARDING PRESCRIBED MEDICATION

Medication shall be administered in school only on a written order by the prescribing physician, along with a written request and a supply of medication from the parent/legal guardian. All medicine must be properly labeled, in the original pharmacy container and brought to school by the parent/legal guardian. Any unauthorized medication found in a student's possession without proper documentation on file, will be taken, held in the school office, and the parent/guardian notified. This is for the safety of your child and others.

Medication in general, according to state law, will be administered or taken under the supervision of the school nurse. Please note that a school nurse is not always available during school hours to administer medication. Receipt of a doctor's order and written request from the parent does not guarantee that a medication can be administered during the school day.

A medication order is effective July 1-June 30 of each school year and $\underline{\text{must}}$ be renewed annually.

In the case of a POTENTIALLY LIFE-THREATENING CONDITION (i.e., epinephrine/inhaler/pancreatic enzymes), legislation has been passed which allows a student to carry a medication for immediate availability and self-administration. However, this situation REQUIRES that you contact the school principal. These medications that may be carried by a student require proper documents to be completed by the student's healthcare provider and parent. In the case of a student with a potentially life-threatening allergy, with documented history of an actual anaphylactic episode, provision of a nurse-trained designee for administration of emergency epinephrine, in the event a nurse is unavailable, is allowable under law. However, certain restrictions apply and you must contact the school nurse.

Sincerely,

Nonpublic Nurse

GLOUCESTER CO. SPECIAL SERVICES SCHOOL DISTRICT NONPUBLIC NURSING PROGRAM 1340 Tanyard Road Sewell, NJ 08080 (856) 468-6530 x1045 ST. MARGARET REGIONAL SCHOOL
773 Third Street
Woodbury Heights, N.J. 08097

Woodbury Heights, NJ 08097 (856) 845-5200 Fax # (856) 845-2405

PHYSICIAN MEDICATION ORDER FORM •SIGNED ORIGINAL ORDER REQUIRED•

Student's Name		Grade	DOB
Nonpublic School			
* PLEASE PROVIDE A SEPAR	ATE FORM FOR EACH	MEDICATION THAT	IS TO BE ADMINISTERED.
*PHYSICIAN TO COMPLET Diagnosis:			
Medication:		DC Dat	te:
Special Instructions:			-
Precautions/Side Effe	ects:		
Date	Physician Signature	(Original / No	
Physician Name			
Address			
Telephone No			
determine the manner in w	urs to administer this r hich medication will b ive July 1 - June 30 of	nedication. Please e dispensed in the each school year a	contact the school principal to absence of a GCSSSD nurse. nd must be renewed annually.
to receive medication at sch			
I WILL BRING THE MEDICA	ATION (PRESCRIPTI ER. PROPERLY LA	ON OR NON-PRE BELED, AND WI	
 Date	Pa	rent/Legal Guardia	an Signature

GLOUCESTER COUNTY SPECIAL SERVICES SCHOOL DISTRICT PERMISSION FOR EMERGENCY ADMINISTRATION OF EPINEPHRINE

THIS ORDER MUST BE RETURNED IN ITS $\underline{\mathbf{ORIGINAL\ FORM}}$. FAXES AND COPIES WILL $\underline{\mathbf{NOT\ BE\ ACCEPTED}}$.

Parent/Guardian Signature		Date
should my child's condition require it. district employee, chief school adn be responsible for any liability as a administration of epinephrine to n	further understand that n nistrator of a nonpublic so esult of any injury arising / child and that I shall in s or agents against any c	ar and must be renewed for each school year, neither the GCSSSD Board of Education, any chool, nor nonpublic school employee shall from the procedures utilized for emergency demnify and hold harmless the district or laims arising out of the administration of a pinephrine to my child.
me) prescribed by our physician/or rof anaphylaxis and does not have the is not available, a designee will ad epinephrine for anaphylaxis to my chithe school nurse using the "Protocol the school the	rrse practitioner for anaphy capability for self-administrationinister a pre-filled, single I. The designee has been plant and Implementation Plant the School Nurse" establists	n for the Emergency Administration of shed by the Department of Education in
(14)	ne of Student)	(Nonpublic School)

Asthma Treatment Plan – Student

(This asthma action plan meets NJ Law N.J.S.A. 18A:40-12.8) (Physician's Orders)







	INE STA
-1	8.32 B
th	D
Health	

(Please Print) Date of Birth Effective Date Name Parent/Guardian (if applicable) **Emergency Contact** Doctor Phone Phone Phone Triggers Check all items Take daily control medicine(s). Some inhalers may be HEALTHY (Green Zone) || || || more effective with a "spacer" - use if directed. that trigger You have all of these: HOW MUCH to take and HOW OFTEN to take it patient's asthma: · Breathing is good 2 puffs twice a day ☐ Advair® HFA ☐ 45, ☐ 115, ☐ 230 ______ ☐ Colds/flu _□ 1, □ 2 puffs twice a dav · No cough or wheeze ☐ Aerospan[™] ☐ Exercise ☐ Alvesco® ☐ 80, ☐ 160 ☐ Dulera® ☐ 100, ☐ 200 ☐ ☐ 1, ☐ 2 puffs twice a day Sleep through □ Allergens _2 puffs twice a day the night o Dust Mites, _2 puffs twice a day ☐ Flovent® ☐ 44, ☐ 110, ☐ 220 _ dust, stuffed · Can work, exercise, ☐ Qvar® ☐ 40, ☐ 80 _ 1, 2 puffs twice a day animals, carpet and play ☐ 1, ☐ 2 puffs twice a day □ Symbicort® □ 80, □ 160 □ o Pollen - trees. ☐ Advair Diskus® ☐ 100, ☐ 250, ☐ 500 ___ _1 inhalation twice a day grass, weeds ☐ Asmanex® Twisthaler® ☐ 110, ☐ 220___ _□ 1, □ 2 inhalations □ once or □ twice a day o Mold ☐ Flovent® Diskus® ☐ 50 ☐ 100 ☐ 250 _____1 inhalation twice a day O Pets - animal _□ 1, □ 2 inhalations □ once or □ twice a day □ Pulmicort Flexhaler® □ 90, □ 180 ___ dander \square Pulmicort Respules® (Budesonide) \square 0.25, \square 0.5, \square 1.0 $\underline{\hspace{0.3cm}}$ 1 unit nebulized \square once or \square twice a day o Pests - rodents, ☐ Singulair® (Montelukast) ☐ 4, ☐ 5, ☐ 10 mg _____1 tablet daily cockroaches □ Other □ Odors (Irritants) O Cigarette smoke ☐ None And/or Peak flow above & second hand Remember to rinse your mouth after taking inhaled medicine. smoke ___ puff(s) ____minutes before exercise. If exercise triggers your asthma, take o Perfumes, cleaning products, CAUTION (Yellow Zone) IIII Continue daily control medicine(s) and ADD quick-relief medicine(s). scented You have any of these: products HOW MUCH to take and HOW OFTEN to take it MEDICINE O Smoke from · Cough burning wood, ☐ Albuterol MDI (Pro-air® or Proventil® or Ventolin®) _2 puffs every 4 hours as needed Mild wheeze inside or outside ☐ Xopenex® ______2 puffs every 4 hours as needed ☐ Albuterol ☐ 1.25, ☐ 2.5 mg ______1 unit nebulized every 4 hours as needed Tight chest ■ Weather o Sudden · Coughing at night __1 unit nebulized every 4 hours as needed temperature ☐ Duoneb® _____ Other: change ☐ Xopenex® (Levalbuterol) ☐ 0.31, ☐ 0.63, ☐ 1.25 mg _1 unit nebulized every 4 hours as needed o Extreme weather ☐ Combivent Respimat® ______1 inhalation 4 times a day - hot and cold If quick-relief medicine does not help within ☐ Increase the dose of, or add: Ozone alert days 15-20 minutes or has been used more than □ Other ☐ Foods: 2 times and symptoms persist, call your If quick-relief medicine is needed more than 2 times a doctor or go to the emergency room. week, except before exercise, then call your doctor. And/or Peak flow from_____ to_ Other: Take these medicines NOW and CALL 911. EMERGENCY (Red Zone) ||||| Your asthma is Asthma can be a life-threatening illness. Do not wait! getting worse fast: HOW MUCH to take and HOW OFTEN to take it MEDICINE Quick-relief medicine did Albuterol MDI (Pro-air® or Proventil® or Ventolin®) 4 puffs every 20 minutes not help within 15-20 minutes 4 puffs every 20 minutes ☐ Xopenex[®] This asthma treatment · Breathing is hard or fast 1 unit nebulized every 20 minutes ☐ Albuterol ☐ 1.25, ☐ 2.5 mg ____ plan is meant to assist. Nose opens wide Ribs show 1 unit nebulized every 20 minutes not replace, the clinical ☐ Duoneb® Trouble walking and talking decision-making ☐ Xopenex® (Levalbuterol) ☐ 0.31, ☐ 0.63, ☐ 1.25 mg ___1 unit nebulized every 20 minutes · Lips blue · Fingernails blue And/or required to meet 1 inhalation 4 times a day ☐ Combivent Respimat® ____ Other: Peak flow individual patient needs. □ Other below Permission to Self-administer Medication: PHYSICIAN/APN/PA SIGNATURE DATE Physician's Orders ☐ This student is capable and has been instructed in the proper method of self-administering of the PARENT/GUARDIAN SIGNATURE non-nebulized inhaled medications named above

REVISED MAY 2017

- in accordance with NJ Law.
- ☐ This student is <u>not</u> approved to self-medicate.

PHYSICIAN STAMP

Make a copy for parent and for physician file, send original to school nurse or child care provider.

Asthma Treatment Plan – Student Parent Instructions

The **PACNJ Asthma Treatment Plan** is designed to help everyone understand the steps necessary for the individual student to achieve the goal of controlled asthma.

- 1. Parents/Guardians: Before taking this form to your Health Care Provider, complete the top left section with:
 - · Child's name
- · Child's doctor's name & phone number
- Parent/Guardian's name

- . Child's date of birth
- An Emergency Contact person's name & phone number
- & phone number

- 2. Your Health Care Provider will complete the following areas:
 - . The effective date of this plan
 - . The medicine information for the Healthy, Caution and Emergency sections
 - Your Health Care Provider will check the box next to the medication and check how much and how often to take it
 - Your Health Care Provider may check "OTHER" and:
 - Write in asthma medications not listed on the form
 - * Write in additional medications that will control your asthma
 - Write in generic medications in place of the name brand on the form
 - Together you and your Health Care Provider will decide what asthma treatment is best for your child to follow
- 3. Parents/Guardians & Health Care Providers together will discuss and then complete the following areas:
 - Child's peak flow range in the Healthy, Caution and Emergency sections on the left side of the form
 - · Child's asthma triggers on the right side of the form
 - <u>Permission to Self-administer Medication</u> section at the bottom of the form: Discuss your child's ability to self-administer the inhaled medications, check the appropriate box, and then both you and your Health Care Provider must sign and date the form
- 4. Parents/Guardians: After completing the form with your Health Care Provider:
 - Make copies of the Asthma Treatment Plan and give the signed original to your child's school nurse or child care provider
 - · Keep a copy easily available at home to help manage your child's asthma
 - Give copies of the Asthma Treatment Plan to everyone who provides care for your child, for example: babysitters, before/after school program staff, coaches, scout leaders

PARENT AUTHORIZATION I hereby give permission for my child to receive medication at schein its original prescription container properly labeled by a pharminformation between the school nurse and my child's health counderstand that this information will be shared with school staff or	nacist or physician. I also giv are provider concerning my	ve permission for the release and exchange of
Parent/Guardian Signature	Phone	Date
FILL OUT THE SECTION BELOW ONLY IF YOUR HEALTH CARE SELF-ADMINISTER ASTHMA MEDICATION ON THE FRONT OF RECOMMENDATIONS ARE EFFECTIVE FOR ONE (1) SCHOOL	THIS FORM.	
☐ I do request that my child be ALLOWED to carry the following in school pursuant to N.J.A.C:.6A:16-2.3. I give permission for replan for the current school year as I consider him/her to be remedication. Medication must be kept in its original prescription shall incur no liability as a result of any condition or injury arise on this form. I indemnify and hold harmless the School District, or lack of administration of this medication by the student.	ny child to self-administer med sponsible and capable of trans on container. I understand tha ing from the self-administrati	sporting, storing and self-administration of the the school district, agents and its employees on by the student of the medication prescribed
\square I D0 N0T request that my child self-administer his/her asthm	na medication.	
Parent/Guardian Signature	Phone	Date



Disclaimers: The use of this Website/PACNJ Ashma Treatment Plan and its content is all your own risk. The content is provided on an "as is" basis. The American Lung Association of the Mid-Altantic (ALAM-A), the Pediatric/Adult Ashma Coalition of New Jersey and all affiliates disclaim all warranties, express or implied, statutory or otherwise, including but not limited to the implied warranties or merchantability, non-infringement of third parties' rights, and filmss for a particular purpose. ALAM-A makes no warranty, representations or warranties about the accuracy, reliability, completeness, currency, or limeliness of the content. ALAM-A makes no warranty, representation or guaranty that the increasing with the increasing without limitability, incidental and consequential damages, personal injury/wronglud death, lost profits, or damages resulting from data or business interruption) resulting from the use or inability to use the content of this Ashma Treatment Plan whether based on warranty, contract, tort or any other legal theory, and whether or not ALAM-A is advised of the possibility of such damages. ALAM-A and its affiliates are not liable for any claim, whatsoever, caused by your use or misuse of the Ashma Treatment Plan, nor of this website.

The Pediatric/Adult Asthma Coalition of New Jersey, sponsored by the American Lung Association in New Jersey. This publication was supported by a grant from the New Jersey Department of Health and Senior Services, with funds provided by the U.S. Centers for Disease Control and Prevention under Cooperative Agreement 5U59EH000491-5. Its content are solely the responsibility of the authors and do not necessarily represent the official views of the New Jersey Department of Health and Senior Services or the U.S. Centers for Disease Control and Prevention. Although this document has been funded wholly or in part by the United States Environmental Protection Agreement Ay66296601-2 to the American Lung Association in New Jersey, although this document has been funded wholly or in part by the United States Environmental Protection Agreement Ay66296601-2 to the American Lung Association in New Jersey, and no official and prevention. Although this document has been funded wholly or in part by the United States Environmental Protection Agreement Ay6629601-2 to the American Lung Association in New Jersey, and no official and environmental Protection Agreement Ay6629601-2 to the American Lung Association in New Jersey, and no official and environmental Protection Agreement Ay6629601-2 to the American Lung Association in New Jersey, and no official and the Protection Agreement Ay6629601-2 to the American Lung Association in New Jersey, and no official and the Protection Agreement Ay6629601-2 to the American Lung Association in New Jersey, and no official and the Protection Agreement Ay6629601-2 to the American Lung Association in New Jersey, and no official and the Protection Agreement Ay6629601-2 to the American Lung Association in New Jersey, and no official and the Protection Agreement Ay6629601-2 to the American Lung Association in New Jersey, and no official Agreement Ay6629601-2 to the American Lung Association in New Jersey, and the Protection Agreement Ay6629601-2 to the American Lung Association in New Jersey, and th



GLOU. CO. SPECIAL SERVICES SCHOOL DISTRICT NONPUBLIC NURSING PROGRAM 1340 Tanyard Road Sewell, NJ 08080 (856) 468-6530 x1045

ST MARGARET REGIONAL SCHOOL 773 Third Street Woodbury Heights, NJ 08097 (856) 845-5200

PHYSICIANS AUTHORIZATION ASSURANCE STATEMENT FOR STUDENT'S SELF-ADMINISTRATION OF MEDICATION

I certify that, a life threatening condition. I am recomm	
self-administer medication. He/She has been instructed by me in the proself-administering the following medication:	ending that the above-named student be permitted to oper method of self-administration, and is capable of
Name and purpose of medication:	
Identification of chronic medical problems:	
Prescribed dosage and schedule:	
Length of time medication is to be taken:	
Possible side effects and/or special precautions:	
Prescribing Physician/Nurse Practioner Signature	Date
Prescribing Physician Murse Practioner Signature	Date
PRINT Physician's/Nurse Practioner's Name	Telephone Number
PARENTAL PERMISSION, RELEASE, INDEMNITY	AND CONSENT TO TREATMENT
and, as p	
ool. I understand that this permission is valid only for this school year d's condition require it. onsideration of the consent and permission for our child to carry and self-ad inst St. Margaret's School, St. Margaret's Church, Woodbury Heights, NJ, the vices School District, and their respective agents, servants, employees, officer	Iminister this medication, we hereby release any and all clain the Diocese of Camden, New Jersey, Gloucester County Species, trustees, administrators, and volunteers, for damages and/
ies to us or our child which may arise from the carrying or use of this medica iduals harmless from and against any claim or claims brought by or on beha in any way connected with our child's possession or use of this medication	alf of our child or by or on behalf of any other person arising of
consent and give our permission for our child to be diagnosed, treated, and/onsed medical personnel. We hereby release any and all claims against St. Mad Diocese of Camden, New Jersey, Gloucester County Special Services School Distees, administrators, and volunteers, for damages and/or injuries to us or outdemnify and hold these entities and individuals harmless from and against a pehalf of any other person arising out of or in any way connected with such consibility as a result of such treatment.	argaret's School, St. Margaret's Church, Woodbury Heights, N strict, and their respective agents, servants, employees, officer or child which may arise from such treatment. We further agre ony claim or claims brought by or on behalf of our child or by
Parent/Guardian Signature	
	Parent/Guardian Signature
PRINT Parent/Guardian Name	Parent/Guardian Signature PRINT Parent/Guardian Name

FUUD ALLEKGY & ANAPHYLAXIS EMERGENCY CARE PL _____ D.O.B.: ____ **PLACE** Name: **PICTURE** Allergic to: HERE _____Ibs. Asthma: ☐ Yes (higher risk for a severe reaction) ☐ No NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE. Extremely reactive to the following allergens: THEREFORE: ☐ If checked, give epinephrine immediately if the allergen was LIKELY eaten, for ANY symptoms. ☐ If checked, give epinephrine immediately if the allergen was DEFINITELY eaten, even if no symptoms are apparent. FOR ANY OF THE FOLLOWING: MILD SYMPTOMS **SEVERE** SYMPTOMS HEART THROAT Itchy or Itchy mouth A few hives, MOUTH runny nose, mild itch nausea or Shortness of Pale or bluish Tight or hoarse Significant swelling of the sneezing discomfort breath, wheezing, skin, faintness, throat, trouble repetitive cough weak pulse, breathing or tongue or lips FOR MILD SYMPTOMS FROM MORE THAN ONE dizziness swallowing SYSTEM AREA, GIVE EPINEPHRINE. OR A FOR MILD SYMPTOMS FROM A SINGLE SYSTEM COMBINATION AREA, FOLLOW THE DIRECTIONS BELOW: of symptoms SKIN from different Repetitive Many hives over Feeling 1. Antihistamines may be given, if ordered by a body areas. vomiting, severe something bad is body, widespread healthcare provider. about to happen, redness diarrhea Stay with the person; alert emergency contacts. anxiety, confusion 3. Watch closely for changes. If symptoms worsen, Û T T give epinephrine. INJECT EPINEPHRINE IMMEDIATELY.

- Call 911. Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
- Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
- Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
- Alert emergency contacts.
- Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.


MEDICATIO	NS/DOS	ES
Epinephrine Brand or Generic:		
Epinephrine Dose: 🗆 0.1 mg IM	□ 0.15 mg IM	□ 0.3 mg IM
Antihistamine Brand or Generic: _		
Antihistamine Dose:		
Other (e.g., inhaler-bronchodilator	if wheezing):	



FUUD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN


HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

- 1. Remove Auvi-Q from the outer case. Pull off red safety guard.
- 2. Place black end of Auvi-Q against the middle of the outer thigh.
- 3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
- 4. Call 911 and get emergency medical help right away.



HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

- Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
- 2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
- 3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
- 4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENACLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

- 1. Remove epinephrine auto-injector from its protective carrying case.
- 2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
- 3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
- 4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.

HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

- 1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
- 2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
- 3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
- 4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
- 5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.

HOW TO USE SYMJEPI™ (EPINEPHRINE INJECTION, USP)

- When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
- 2. Hold SYMJEPI by finger grips only and slowly insert the needle into the thigh. SYMJEPI can be injected through clothing if necessary.
- 3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
- 4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
- 5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.

ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

- 1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
- 2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
- 3. Epinephrine can be injected through clothing if needed.
- 4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

EMERGENCY CONTACTS — CALL 911
OTHER EMERGENCY CONTACTS

RESCUE SQUAD:
NAME/RELATIONSHIP:
PHONE:

DOCTOR:
PHONE:
NAME/RELATIONSHIP:
PHONE:

PARENT/GUARDIAN:
PHONE:
NAME/RELATIONSHIP:
PHONE:

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

	2				390
•	J	£	1	0.00	
		4	1/1	1	
		ŀ		THE CONTRACTOR	d
400					

FORM PROVIDED COLIRTESY OF FOOD ALLERGY RESEARCH & EDUCATION (FARE) (FOODALLERGY ORG) 5/2020